

Total No. of Questions—8]

[Total No. of Printed Pages—4+1

Seat No.	

[5667]-1001

F.E. (I Semester) EXAMINATION, 2019

ENGINEERING MATHEMATICS—I

(Phase-II)

(2019 PATTERN)

Time : 2½ Hours

Maximum Marks : 70

N.B. — (i) Attempt Q. No. 1 or Q. No. 2, Q. No. 3 or Q. No. 4, Q. No. 5 or Q. No. 6, Q. No. 7 or Q. No. 8.

- (ii) Use of electronic pocket calculator is allowed.
- (iii) Assume suitable data, if necessary.
- (iv) Neat diagrams must be drawn wherever necessary.
- (v) Figures to the right indicate full marks.

1. (a) If $z = \tan(y + ax) + (y - ax)^{3/2}$, find the value of

$$\frac{\partial^2 z}{\partial x^2} - a^2 \frac{\partial^2 z}{\partial y^2}.$$

[6]

(b) If $T = \sin\left(\frac{xy}{x^2 + y^2}\right) + \sqrt{x^2 + y^2}$, by using Euler's theorem

$$\text{find } x \frac{\partial T}{\partial x} + y \frac{\partial T}{\partial y}.$$

[6]

(c) If $u = x^2 - y^2$, $v = 2xy$ and $z = f(u, v)$, then show that

$$x \frac{\partial z}{\partial x} - y \frac{\partial z}{\partial y} = 2\sqrt{u^2 + v^2} \frac{\partial z}{\partial u}.$$

[6]

P.T.O.

Or

2. (a) If $x = u \tan v$, $y = u \sec v$, prove that : [6]

$$\left(\frac{\partial u}{\partial x} \right)_y \left(\frac{\partial v}{\partial x} \right)_y = \left(\frac{\partial u}{\partial y} \right)_x \left(\frac{\partial v}{\partial y} \right)_x.$$

- (b) If $u = \sin^{-1} \left(\frac{x+y}{\sqrt{x+y}} \right)$, by using Euler's theorem.

prove that : [6]

$$x^2 \frac{\partial^2 u}{\partial x^2} + 2xy \frac{\partial^2 u}{\partial x \partial y} + y^2 \frac{\partial^2 u}{\partial y^2} = \frac{1}{4} (\tan^3 u - \tan u).$$

- (c) If $x = \frac{\cos \theta}{u}$, $y = \frac{\sin \theta}{u}$ and $z = f(x, y)$, then show that : [6]

$$u \frac{\partial z}{\partial u} - \frac{\partial z}{\partial \theta} = (y-x) \frac{\partial z}{\partial x} - (y+x) \frac{\partial z}{\partial y}.$$

3. (a) If $u = x + y + z$, $v = x^2 + y^2 + z^2$, $w = xy + yz + zx$

find $\frac{\partial(u, v, w)}{\partial(x, y, z)}$. [6]

- (b) Examine whether $u = \frac{x-y}{1+xy}$, $v = \tan^{-1} x - \tan^{-1} y$ are functionally dependent, if so find the relation between them. [5]

- (c) Find the extreme values of $x^2 + y^2 + \frac{2}{x} + \frac{2}{y}$. [6]

Or

4. (a) If $u = x + y^2$, $v = y + z^2$, $w = z + x^2$, using Jacobian find $\frac{\partial x}{\partial u}$. [6]

- (b) A power dissipated in a resistor is given by $P = \frac{\varepsilon^2}{R}$. If errors of 3% and 2% are found in ε and R respectively, find the percentage error in P . [5]
- (c) Using Lagrange's method find extreme value of xyz if $x + y + z = a$. [6]

5. (a) Examine for consistency of the system of linear equations and solve if consistent : [6]

$$\begin{aligned}x_1 + x_2 + x_3 &= 0 \\ -2x_1 + 5x_2 + 2x_3 &= 1 \\ 8x_1 + x_2 + 4x_3 &= -1\end{aligned}$$

- (b) Examine for linear dependence or independence the vectors $(1, 1, 1, 3)$, $(1, 2, 3, 4)$, $(2, 3, 4, 7)$. Find the relation between them if dependent. [6]
- (c) Determine the values of a , b , c when A is orthogonal where : [5]

$$A = \begin{bmatrix} 0 & 2b & c \\ a & b & -c \\ a & -b & c \end{bmatrix}.$$

Or


6. (a) Investigate for what values of a and b , the system of equations $2x - y + 3z = 2$, $x + y + 2z = 2$, $5x - y + az = b$ have :
- (1) No solution
 - (2) A unique solution
 - (3) An infinite number of solutions. [6]

(b) Examine for linear dependence or independence the vectors

$$x_1 = (2, 3, 4, -2), x_2 = (1, 1, 2, -1), x_3 = \left(\frac{-1}{2}, -1, -1, \frac{1}{2}\right).$$

Find the relation between them if dependent. [6]

(c) Determine the currents in the network given in figure below : [5]



7. (a) Find the eigen values and the corresponding eigen vectors for the following matrix : [6]

$$A = \begin{bmatrix} 4 & 0 & 1 \\ -2 & 1 & 0 \\ -2 & 0 & 1 \end{bmatrix}.$$

(b) Verify Cayley-Hemilton theorem for $A = \begin{bmatrix} 1 & -1 & 0 \\ 2 & 3 & -2 \\ -2 & 0 & 1 \end{bmatrix}$ and use it to find A^{-1} . [6]

(c) Find a matrix P that diagonalizes the matrix

$$A = \begin{bmatrix} 1 & 6 & 1 \\ 1 & 2 & 0 \\ 0 & 0 & 3 \end{bmatrix}. [6]$$

Or

8. (a) Find the eigen values and the corresponding eigen vectors for the following matrix : [6]

$$A = \begin{bmatrix} 5 & 0 & 1 \\ 0 & -2 & 0 \\ 1 & 0 & 5 \end{bmatrix}.$$

- (b) Verify Cayley-Hamilton theorem for $A = \begin{bmatrix} 0 & 1 & 0 \\ 0 & 0 & 1 \\ 1 & -3 & 3 \end{bmatrix}$ and use it to find A^{-1} . [6]

- (c) Reduce the following quadratic form to the sum of the squares form : [6]

$$Q = 2x^2 + 9y^2 + 6z^2 + 8xy + 8yz + 6xz.$$